

Microsporium canis and other dermatophytes isolated from humans, dogs and cats in Mexico city

Torres-Guerrero E¹, González de Cossío AC¹, Segundo ZC², Cervantes ORA³, Ruiz-Esmenjaud J¹ and Arenas R¹

¹Sección de Micología. Hospital “Dr. Manuel Gea González”, Ciudad de México, México

²Laboratorio de Micología Veterinaria, Unidad de Servicios de Diagnóstico y Constatación del Centro de Enseñanza, Investigación y Extensión en Producción Animal en Altiplano. FMVZ.UNAM, México

³Laboratorio de Micología del Depto. de Microbiología e Inmunología de la FMVZ. UNAM. México

Abstract

Introduction: Tinea or ringworm are infections caused by dermatophytes, a group of keratinophilic fungi.

According to origin and tropism can be classified as anthropophilic, zoophilic and geophilic.

Objectives: To obtain current data in zoophilic dermatophytes in humans and animals in Mexico City.

Material and methods: We performed an observational, descriptive and retrospective study, in two different Departments of Mycology. In a General Hospital we studied dermatophytic infections in 8,684 patients, (human cases), and in 480 animals (377 dogs and 103 cats) at the Faculty of Medical Veterinary, University of Mexico.

Results: 57 of the 8,684 human patients (0.65%), had zoophilic dermatophytes: *M. canis* 43 (75.5%), *T. mentagrophytes var. mentagrophytes* 13 (22.9%) and *M. nanum* 1 (1.6%); 40.8% were men and 59.2% females.

At the Veterinary Faculty, 377 samples from dogs and 103 from cats were studied, with 33 (8.73 %) and 36 (34.95%) positive cultures respectively. In dogs *M. canis* 72.70% *Trichophyton terrestris* 12.15%, *M. gypsum* 9.10% and *T. mentagrophytes* 6.10%, and in cats only *M. canis* was isolated.

Conclusions: In humans and in animals *M. canis* is still the main causal zoophilic agent. In children tinea capitis is the most frequent dermatophytosis. Tinea corporis is the second one, and usually related to close contact with infected pets.

Introduction

Tinea or ringworm are infections caused by keratinophilic fungi called dermatophytes that can also invade the skin and its appendages [1]. There are three anamorphic genders: *Trichophyton*, *Epidermophyton* and *Microsporium*, none of which form part of the cutaneous flora. They can be classified according to its origin and tropism in anthropophilic, zoophilic and geophilic. These infections constitute 70 to 80% of all the mycoses and represent 5% of the dermatological consults [2,3].

Infection is by direct contact with the causal agent and it can appear in any race, sex, socioeconomic level or occupation [3]. The animals act as reservoirs and can be symptomatic or just be carriers.

Among the zoophilic dermatophytes, *Microsporium canis* and *Trichophyton mentagrophytes*, are of main medical relevance in Mexico.

The tinea are frequent in domestic and savage animals; they are found in bovine, pigs and equine as well as in poultry, the most affected ones, are the small species, such as dogs, cats and rodents. To acquire infection, a direct contact with the contaminated source is needed, soil or animal or it can also be transmitted from person to person or by fomites [3].

Our aim is to obtain the current frequency of the zoophilic dermatophytes in samples collected from patients and animals with a clinical diagnosis of tinea in a general hospital and in a veterinary clinic respectively.

Material and methods

We performed an observational, descriptive and retrospective study, in two departments of mycology, at “Dr. Manuel Gea Gonzalez” General Hospital, and at the Faculty of Medical Veterinary, Autonomous National University of Mexico (UNAM).

During a 10 year-period, a mycological study was performed, in 8684 patients with cutaneous lesions suggestive of tinea in the hospital and in the same period, 480 animals at the Faculty of Medical Veterinary, 377 dogs and 103 cats with suspected dermatophytes. All of them were included.

A direct exam with 20% potassium hydroxide (KOH) and a culture in Sabouraud dextrose agar with cycloheximide and chloramphenicol at 30°C were performed in samples from humans and animals. Identification was performed based in the morphologic criteria by the microscopic observation with lactophenol cotton blue.

Correspondence to: Dr. Roberto Arenas. Sección de Micología. División de Investigación. Hospital “Dr. Manuel Gea González”. Calzada de Tlalpan 4800, Colonia “Belisario Domínguez, sección XVI”, Ciudad de México C.P. 14080. E-mail: rarenas98@hotmail.com

Received: March 01, 2016; **Accepted:** March 31, 2016; **Published:** April 04, 2016

Results

From the 8,684 human patients, zoophilic dermatophytes were isolated only in 57 (0.65%) patients: *M. canis* 43 (75.5%), *T. mentagrophytes* (0.65%) var. *mentagrophytes* 13 (22.9%) and *M. nanum* 1 (1.6%) (Graph 1), 40.8% were men and 59.2% females.

The most affected age groups were 0 to 10 (46.89%); and the group 21 to 30 (12.5%) (Graph 2). Clinical features are shown in Graph 3.


At the Veterinary Faculty, 480 samples from dogs and cats were studied: 377 from dogs and 103 from cats, with 33 (8.73%) and 36 (34.95%) positive cultures respectively. In dogs *M. canis* (72.70%), *Trichophyton terrestris* (12.15%), *M. gypseum* (9.10%) and *T. mentagrophytes* (6.10%), were isolated, while in cats the only dermatophyte was *M. canis* (Graph 1).

Discussion


According with 4th National Consensus of Superficial Mycoses in Mexico (2008), 70 to 80% of mycoses in outpatients were caused by dermatophytes [4]: 2% tineamannus, 4% tineacuris, 4-10% tineacapitis, 15% tineacorporis, 30% tineangium and 45% tineapedis [4]. These data are considered on the basis of a general population, but only tineacapitis is almost always present just in children. [5]

In the current study, the majority of the cases in humans correspond to tineacapitis (46.89%) followed by tineacorporis (27.86%) because the highest percentage of patients was found from 0-10 years of age (46.89%). This data is congruent, as tineacapitis was the most frequent affected was the pediatric group, with an exceptional case in a 97 year-old female [5].


M. canis was the most frequent dermatophyte. It is present in 4.1% among general statistics and is the causal agent in 60 to 89% of tineacapitis in Mexican children [3,6]. The contact with domestic animals represents the main source of infection in up to 83%. [3,6] Similar results by Monteagudo [7], after his study conducted in Santiago de Compostela with 196 cases of tineacapitis, observed *M. canis* in 70-95%, and *T. mentagrophytes* var. *mentagrophytes* in 14.8%, and a low frequency of other non-zoophilic species. These data are different from USA reports, where the main causal agent of tineacapitis is *T. tonsurans* (90%), the increase incidence of this agent is related to migrations of Afro-Americans and Latin American [3,8]. Also in Madrid, *T. tonsurans*' incidence has increased in the last years, and *M. canis* is now in the second place. In Puerto Rico about two thirds of tineacapitis are caused by *T. tonsurans* and the other third by *M. canis*



Graph 1. Zoophilic dermatophytes isolated from humans, dogs and cats.



Graph 2. Frequency of dermatophytoses according to age group.



Graph 3. Clinical human dermatophytic infections caused by zoophilic dermatophytes.

[9].

In Jordan after conducting a 6 years epidemiological study, Shagra [10] found *T. violaceum* as the main causal agent in *tineacapitis*, and in second place, *M. canis*. Mseddi and Makni in Tunez, separately reported 83 and 68% *tineacapitis* caused by *T. violaceum*, and *M. canis* in 29.2% [11,12]. Also Ouidaina, in Morocco, after finishing a study with 1299 patients from 1993 to 2007, identified 76.4% of *tineacapitis* due to *T. mentagrophytes*, while *M. canis* 13.4% [13]. In the same country, Boumhil, studied 162 patients from 2002 to 2008, finding 63.58% of infections due to *T. violaceum* and 33.33% to *M. canis* [14]. Arenas *et al.*, reported in an epidemiological study conducted in the urban and rural zones of Dominican Republic, a resurging of *M. audouinii* and *T. tonsurans* respectively, probably due to the migratory movements from Haiti, while *M. canis* was found in 19.04% and 11.65% respectively [15].

Tineacorporis, ranked second in frequency of the clinical forms observed in our study (27.86%). According to the National Consensus of Superficial Mycoses [4], *T. rubrum* was the main causal agent, followed by *M. canis*, however it must be considered that *tineacorporis* caused by *M. canis* is more typical of pediatric patients, as we are reporting in this paper. In a retrospective study in Mexico 357 cases of *tineacorporis* were reported in 21 years, and *M. canis* was found in 16.7% [16]. Meanwhile, Cafarchia *et al.* [17], in 2005, performed in Bari, Italy, a dermatophytic search in dogs and cats, which owner had or had not tinea and found *M. canis* in 53.6% of cats and 36.4% in dogs whose owners had tinea, and in only 14.6% of cats and non in the dogs which owners did not have tinea, concluding that one must be aware that the animals are a source of infection.

Concerning tinea faciei (20% of tinea corporis) and onychomycosis, in this study each one was present in 9.5%. In Italy, Monod performed a study of facial tinea caused by *M. canis*, reporting 91.3% of the children infected by their pets. [18] According to Romano *et al.*, the medium age in tinea faciei is close to 27 years of age, while in Aste *et al.*, paper, it is more common between 36 and 45 years of age. [19] Our facial cases were found in the age group of 20 to 50 years.

We found a lower frequency of *M. canis* in other dermatophytic infections, 9.5% in onychomycoses and 6.25% in tinea pedis. Kazemi, in Iran, studied 590 patients from 1996 till 2004, found zoophilic dermatophytes such as *T. mentagrophytes* and *M. canis*, with a frequency of 65.5% of cases with onychomycosis; Sei in Japan, identified *M. canis* in just 5 cases in 36,052 ambulatory patients. In feet and nails, *M. canis* is uncommon but *Trichophyton* spp are more contagious and responsible of outbreaks in endemic countries, and zoophilic fungi are just related to small familiar epidemics [20-22].

From the veterinary point of view, *Microsporium canis* is the most common dermatophyte in pets [18]. Worldwide 90-100% of dermatophytoses in cats are caused by *M. canis* [23] and usually its isolation in asymptomatic animals indicates subclinical infection or carrier [24].

In the laboratory of the faculty of veterinary medicine at the UNAM, 100% of *M. canis* was reported in the hair of cats and 72.70% in dogs, similar to Lorio in Italy, who showed that the stray cats are an important source of dermatomycosis, because he isolated fungi in 100% of the hair samples [25]. Boumhil in Morocco reported that 56.7% of the cases with tinea capitis have been originated from direct contact with animals [14]. Cafarchia in Italy and Seker in Turkey [26,27], described that in animals one of the main risks to acquire dermatophytoses was the age. Moriello, found that cats with the immunodeficiency Feline virus, the dermatophytosis are three times more prevalent [28].

In this study *M. gypseum* was isolated in a 9.10% of dogs' samples, which differs from the reported percentage by Álvarez *et al.*, in Colombia, who isolated *M. gypseum* in 55.9% in stray dogs and owned dogs in a study in 2001 [29]; Madrid *et al.*, in Brazil, from 7 dogs, 20 day-old, *M. gypseum* was isolated [30].

The isolation of *M. gypseum* in animals, occurs less frequently and its presence is in general attributed to its geophilic character, besides external and internal factors that can alter the normal microbiota of normal animal's skin and this way the infection process is favor, as has been proven in previous studies by Levy *et al.*. Those who determined the presence of dermatophytes in the hair of cats in captivity, found *M. gypseum* in 1.6% of animals and they attributed this finding to the humid conditions, pH and fecal contamination of the place where they are housed [31].

As mentioned before *T. mentagrophytes*, is isolated in different inflammatory entities in the head and nails, but in animals this dermatophyte was isolated just in dogs (6.10%). This dermatophyte has been reported in rodents and less frequently in dogs and other mammals. Murmu *et al.* in a previous study isolated in 16.7% from dogs' hair, skin and nails with suggestive lesions of dermatophytoses [32]. In India Gangil *et al.*, isolated *T. mentagrophytes* in 18.3% from dogs with skin lesions [33], while in Baghdad, Jasim isolated 30.95% in dogs, from samples taken from various body sites [34]. These data contrast with our findings as we found a lower frequency than the previously cited authors, but this finding becomes relevant as it can be a source of infection or re-infection in human and animals, so better

hygienic conditions are recommended to dogs' owners as well as in places where animals are too close to each other (aesthetic canines and veterinary clinics).

T. terrestre was isolated just in dogs (12.15%), similar to *M. gypseum*, though its presence can be attributed to its geophilic character, but different from the last one, it is considered as a causal agent of dermatophytoses in animals. *T. terrestre* does not count with conclusive evidence to confirm its role in dermatophytoses in dogs, attributing their presence to the rather natural behavior of the animal, as when they smell, their snout is in direct contact with the soil, the source of dermatophyte infection [35].

Conclusions

M. canis is the most frequent zoophilic dermatophyte isolated from humans and animals. It is still the main causal dermatophyte for tinea capitis in children and tinea corporis in young adults. This group is the most frequently affected, because afterward in puberty long-chain fat acids are in the scalp and have a protective role.

Also, children are more often in direct contact with pets. Infections may be related with poor hygiene especially when animals are too close to each other (aesthetics and veterinary clinics) and can become a focus of infection. Zoophilic or geophilic dermatophytes could also be the cause of family epidemics.

References

- López-Martínez R, Méndez-Tovar LJ, Hernández-Hernández F, Castañón - Olivares R (2006) Micología Médica. Procedimientos para el diagnóstico de laboratorio (2nd edn). En: Trillas, México, 31-46.
- Arenas R (2002) [Dermatophytoses in Mexico]. *Rev Iberoam Micol* 19: 63-67. [Crossref]
- Arenas R (2015) Micología Médica Ilustrada (5th Edn). Mc Graw Hill Companies, México. Pp: 67-98.
- Micosis Superficiales (2008) Cuarta revisión del Consenso Nacional de Prevención, Diagnóstico y Tratamiento. México, Universidad Nacional Autónoma de México. Pp. 20 - 24.
- Torres-Guerrero E, Leal- Osuna S, Clavellina M, Solís P, Arenas R (2014) Tinea capitis en pacientes geriátricas. Informe de dos casos por *Trichophyton tonsurans* y *Microsporium canis*. *Revista Colombiana de Gerontología y Geriatria* 28: 1942-1954.
- Segundo C, Martínez A, Arenas R, Fernández R, Cervantes RA (2004) [Superficial infections caused by *Microsporium canis* in humans and animals]. *Rev Iberoam Micol* 21: 39-41. [Crossref]
- Moriello K (2012) Dermatophytosis (4th edn). Infectious diseases of the dog and cat. Elsevier, St. Louis, USA. Pp: 588-602.
- Elewski B, Hughey L, Sobera OJ, Hay R (2012) Fungal Diseases. Dermatology (3rd Edn). Elsevier Saunders, New York. Pp: 1251-1285.
- Ginter-Hanselmayer G, Weger W, Ilkit M, Smolle J (2007) Epidemiology of tinea capitis in Europe: current state and changing patterns. *Mycoses* 50 Suppl 2: 6-13. [Crossref]
- Abu Shaqra QM, Al Momani W (2011) Cases of tinea capitis as encountered in a private practice laboratory from Jordan. *J Mycol Med* 21: 24-27. [Crossref]
- Makni F, Néji S, Sellami A, Cheikrouhou F, Sellami H, *et al.* (2008) Les teignes du cuir chevelu dans la région de Sfax (Tunisie). *J Med Mycol* 18: 162-165.
- Mseddi M, Merrekchi S, Sellami H, Mnif E, Boudaya S, *et al.* (2005) Les teignes de l'adulte: étude rétrospective dans le sud Tunisien. *J Med Mycol* 15: 93-96.
- Oudaina W, Biougnach H, Riane S, El Yaagoubil I, Tangi R, *et al.* (2011) [Epidemiology of tinea capitis in outpatients at the Children's Hospital in Rabat (Morocco)]. *J Mycol Med* 21: 1-5. [Crossref]
- Boumhil L, Hjira N, Naoui H, Zerroua A, Bhirich N, *et al.* (2010) Les teignes du cuir chevelu à l'hôpital militaire d'instruction Mohammed V (Maroc). *J Med Mycol* 20: 97-100.

15. Arenas R, Torres E, Amaya M, Rivera ER, Espinal A, *et al.* (2010) Tinea capitis. Emergencia de *Microsporum audouinii* y *Trichophyton tonsurans* en la República Dominicana. *Actas Dermosifilogr* 101: 330-335.
16. Guevara–Cervantes JF, Marioni–Manríquez S, Tello–Ibáñez OO, Vega DC, Vázquez del Mercado E, *et al.* (2015) Tinea corporis. Estudio micológico y epidemiológico de 357 casos. *DCMQ* 13: 282-288.
17. Cafarchia C, Romito D, Capelli G, Guillot J, Otranto D (2006) Isolation of *Microsporum canis* from the hair coat of pet dogs and cats belonging to owners diagnosed with *M. canis* tinea corporis. *Vet Dermatol* 17: 327-331. [[Crossref](#)]
18. Monod M (2008) Secreted proteases from dermatophytes. *Mycopathologia* 166: 285-294. [[Crossref](#)]
19. Torres–Guerrero E, Ramos–Betancourt L, Martínez–Herrera E, Arroyo–Camarena S, Porras C, *et al.* (2015) Dermatophytic blefaritis due to *Microsporum gypseum*. An adult variety of tinea faciei with dermatophytoma. *Our Dermatol Online* 6: 36-38.
20. Kazemi A (2007) Tinea unguium in the north-west of Iran (1996-2004). *Rev Iberoam Micol* 24: 113-117. [[Crossref](#)]
21. Sei Y (2015) [2011 Epidemiological Survey of Dermatomycoses in Japan]. *Med Mycol J* 56: J129-135. [[Crossref](#)]
22. Abd Elmegeed AS, Ouf SA, Moussa TA, Eltahlawi SM (2015) Dermatophytes and other associated fungi in patients attending to some hospitals in Egypt. *Braz J Microbiol* 46: 799-805. [[Crossref](#)]
23. Monteagudo B, Pereiro M Jr, Peteiro C, Toribio J (2003) Tinea capitis en el área sanitaria de Santiago de Compostela. *Actas Dermosifilogr* 94: 598-602.
24. Frymus T, Gruffydd-Jones T, Pennisi MG, Addie D, Belák S, *et al.* (2013) Dermatophytosis in cats: ABCD guidelines on prevention and management. *J Feline Med Surg* 15: 598-604. [[Crossref](#)]
25. Iorio R, Cafarchia C, Capelli G, Fasciocco D, Otranto D, *et al.* (2007) Dermatophytoses in cats and humans in central Italy: epidemiological aspects. *Mycoses* 50: 491-495. [[Crossref](#)]
26. Cafarchia C, Romito D, Sasanelli M, Lia R, Capelli G, *et al.* (2004) The epidemiology of canine and feline dermatophytoses in southern Italy. *Mycoses* 47: 508-513. [[Crossref](#)]
27. Seker E, Dogan N (2011) Isolation of dermatophytes from dogs and cats with suspected dermatophytosis in Western Turkey. *Prev Vet Med* 98: 46-51. [[Crossref](#)]
28. Moriello KA (2004) Treatment of dermatophytosis in dogs and cats: review of published studies. *Vet Dermatol* 15: 99-107. [[Crossref](#)]
29. Álvarez MI, Caicedo LD (2001) Dermatofitos en perros de Cali, Colombia. *Biomédica* 21: 128-133.
30. Madrid IM, Dos Reis Gomes A, Souza Mattei A, Santin R, Brum-Cleff M, *et al.* (2012) Dermatofitose neonatal canina por *Microsporum gypseum*. *Vet Zootec* 19: 073-078.
31. Levy Bentubo HD, Luzes Fedullo JD, Ramiro Corrêa SH, Hidalgo R, Teixeira F, *et al.* (2006) Isolation of *Microsporum gypseum* from the haircoat of health wild felids kept in captivity in Brazil. *Braz J Microbiol* 37: 148-152.
32. Murmu S, Debnath C, Pramanik AK, Mitra T, Jana S, *et al.* (2015) Detection and characterization of zoonotic dermatophytes from dogs and cats in and around Kolkata. *Vet World* 8: 1078-1082.
33. Gangil R, Dutta P, Tripathi R, Singathia R, Lakhotia RL (2012) Incidence of dermatophytosis in canine cases presented at Apollo Veterinary College, Rajasthan, India. *Vet World* 5: 682-684.
34. Mohammed JS (2013) Dermatophytes for isolated from dogs suspected of dermatophytosis in Baghdad City. *Diyala Journal for Pure Sciences* 9: 61-66.
35. Viguie-Vallanet C, Paugam A (2009) Dermatofitos transmitidos por animales. *Acta bioquím clin Latinoam* 43: 263-270.